

BLUE SPECTRAL INFLATION

FRANZ E. SCHUNCK¹ AND ECKEHARD W. MIELKE²

¹*Institut für Theoretische Physik, Universität zu Köln, 50923 Köln, Germany*
E-mail: fs@thp.uni-koeln.de

²*Departamento de Física, Universidad Autónoma Metropolitana-Iztapalapa, Apartado Postal 55-534, C.P. 09340, México, D.F., MEXICO*
E-mail: ekke@xanum.uam.mx

The next-to leading order in the slow-roll approximation of *inflation* is known¹ to lead to the nonlinear equation

$$2C\epsilon y^2\epsilon'' - (3\epsilon + 1)y\epsilon' + \epsilon^2 + \epsilon + \Delta = 0, \quad (1)$$

of Stewart and Lyth², where the energy density $\epsilon = \epsilon(y)$ depends on $y = H^2$, the squared Hubble expansion rate, $C := -2 + \ln 2 + \gamma \simeq -0.73$, and $\Delta := (n_s - 1)/2$ denotes the deviation from the scale invariant Harrison–Zel’dovich spectrum.

It is gratifying that some of our new *discrete eigenvalues* of the spectral index in the blue regime $n_s > 1$ have consistent fits to the cumulative COBE data as well as to recent ground-base CMB experiments.

In general, *exact* inflationary solutions, after an elegant coordinate transformation³, depend on the Hubble expansion rate H as a new “inverse time”, and the regime of inflationary potentials allowing a graceful exit has already been classified⁴. In our phenomenological approach, the inflationary dynamics is not prescribed by one’s theoretical prejudice. On the contrary, in this solvable framework, the ‘graceful exit function’ $g(H)$, which determines the *inflaton potential* $U(\phi)$ and the so-called graceful exit to the Friedmann cosmos, is *reconstructed*⁵ in order to fit the data.

Table 1. *Eigenvalues of the spectral index n_s*

Eigenvalue n_s	Spectrum	Potential $U(\phi)$
(−6.57797)	(unrealistic)	$\sim \tan^2(\sqrt{\kappa}\phi/4)$
$-3 < n_s < 1$	continuous	$\propto \exp(\pm\sqrt{2\kappa}A_0\phi)$
$\vdots$	discrete	$\vdots$
1.106	($m = 5$)	
1.125	All CMB data	
1.153	($m = 3$)	
1.194	COBE data	cf. Fig. 2 in ¹
1.247	($m = 1$)	
1.49575		$\sim \tan^2(\sqrt{\kappa}\phi/4)$

Recent astronomical observations by COBE of the cosmic microwave background (CMB) confirm that the Universe expands rather homogeneously on the

large scale. Four years CMB of cumulative observations, have yielded $n_s = 1.2 \pm 0.3$ as spectral index, including the quadrupole ($l = 2$) anisotropy. This experiment has been complemented by ground-based CMB experiments. Nowadays best-fit to *all* CMB data⁶ is a Hubble constant of $H_0 = 65 \text{ km/(s Mpc)}$ and requires *dark energy* with $\Omega_\Lambda = 0.69$ and a spectral index $n_s = 1.12$.

We have applied the H -formalism³ to the rather accurate first order corrections to the slow-roll approximation of Stewart and Lyth, and transform their nonlinear equation into the *Abel equation* (1). The spectral index of our new class¹ of solutions can only adopt the following *discrete eigenvalues*

$$n_s = 1 + \frac{(2m-1)[4m-1-4m(2m-1)C]}{2[2m(2m-1)C-3m+1]^2} \simeq 1 - \frac{1}{2mC}, \quad (2)$$

where, for large $m \geq 1$, we find asymptotically the scale invariant Harrison-Zel'dovich solution with index $n_s = 1$ as *limiting point*. Let us stress that our discrete spectrum approaches it from the *blue* side, which previously was considered rather difficult to achieve. In order to correlate this with observational restrictions, we display the highest eigenvalues for n_s in Table 1. For the new *blue* spectrum (2) we have reconstructed for $m = 1$ the potential $U(\phi)$ numerically in Fig. 1 of Ref.¹.

Recently, there have been raised some doubts⁷ on the applicability of the second order slow-roll approximation. More recently, however, Stewart and Gong⁸ have explicitly calculated the spectral index up to *second-order corrections*, thereby also corroborating their previous next-to leading order approximation. Then the Abel equation (1) turns into a third order nonlinear differential equation for the energy density with $(2C - \pi^2/6)\epsilon^2 y^3 \epsilon'''$ as leading term. While our solutions of (1) are derived from an *even* Taylor expansion in $y = H^2$, the form of the next order corrections suggests that the additional terms will be *odd* in y . Consequently, our reconstruction on the basis of the Abel equation is maintained to some approximation, only 'orthogonal' terms are expected to arise from second order corrections on the percent level.

References

1. F.E. Schunck and E.W. Mielke, *Phys. Lett. B* **485**, 231 (2000).
2. E.D. Stewart and D.H. Lyth, *Phys. Lett. B* **302**, 171 (1993).
3. F.E. Schunck and E.W. Mielke, *Phys. Rev. D* **50**, 4794 (1994).
4. E.W. Mielke and F.E. Schunck, *Phys. Rev. D* **52**, 672 (1995).
5. J.E. Lidsey, A.R. Liddle, E.W. Kolb, E.J. Copeland, T. Barriero and M. Abney, *Rev. Mod. Phys.* **69**, 373 (1997).
6. S. Dodelson and L. Knox, *Phys. Rev. Lett.* **84**, 3523 (2000).
7. J. Martin and D. Schwarz, *Phys. Rev. D* **62**, 103520 (2000).
8. E.D. Stewart and J.-O. Gong: "The density perturbation power spectrum to second-order corrections in the slow-roll expansion", astro-ph/0101225 v2.